



Taurolidine-Citrate-Heparin the superior lock solution

Clinical Evidence
for central venous catheters in haemodialysis



Introduction

Catheter-related infections and dysfunction are the main catheter complications causing morbidity and mortality in haemodialysis patients. Over the years, many lock solutions - often based on sodium citrate - have been discussed as best alternative to 5.000 IU/mL Heparin. Suppliers of 4% Citrate containing lock solutions even claim it is antimicrobial. However, there is no supporting clinical evidence.



Expert Recommendations

According to the European Renal Best Practice (ERBP 2010), in two studies (McRae, Grudzinski), no benefit regarding infectious complications had been observed for 4% citrate.^{2,3}

4% Citrate vs Heparin ³		
	Heparin 5000 IU/ml	4% sodium citrate
CRBSI rate per 1.000 catheter days	3.3	2.2
Not significant		


Antibacterial lock solutions should ... be preferred. ... taurolidine-citrate solutions can be used.⁴

Taurolidine-Citrate vs Heparin ⁴		
	Heparin 5000 IU/ml	Taurolidine + 4% Sodium Citrate
CRBSI rate per 1.000 catheter days	5.6	decreased by about 90%
Bacteremia-free survival at 90 days (p<0.001)	47%	94%

“

Allon recommended in 2003: Because of a number of incidents of clotting, the possibility of adding Heparin to the Taurolidine/Citrate as soon as the catheter is inserted, should be investigated.

”⁵



Taurolidine - the solution is in the combination


In 2012: Solomon confirmed „...the addition of 500 IU/ml of Heparin to Taurolidine-Citrate may achieve patency comparable to Heparin 5000 U/ml.“⁶

“
**The most promising results are obtained when
antibacterial compounds are added to citrate or heparin.**
”⁷



TauroLOCK™ HEP 500

Taurolock™-Hep500 contains the antimicrobial compound Taurolidine, 4% sodium citrate and 500 IU/mL Heparin.



Additional benefits of Urokinase

The utilisation of TauroLock™-Hep500 as an **antimicrobial** lock solution in dialysis has shown a significant reduction of catheter related bloodstream infections (CRBSI) in many different studies.^{8,9,10}

Even better results are achieved with the so called **2+1 protocol**:

Twice per week TauroLock™-Hep500 and once per week TauroLock™-U25.000.

		TauroLock™ HEP 500	2+1 protocol TauroLock™ HEP 500 TauroLock™ U25.000
Patency	Total number of CVC exchanges	7	1
	rt-PA Rate per 1.000 catheter days	4.3	2.1
CRBSI	per 1.000 catheter days	0.3	0.08


 Study authors: Al-Ali, F. at al.¹¹

 Study location: Qatar  Patients enrolled: 164

 Journal and publication date: Nephrology Dialysis Transplantation (ndt), 2017

 Keynotes: "Taurolock™-U25.000 is a safe and effective tunnelled dialysis catheter lock solution, with low rate of catheter exchange."

"No reported serious adverse event or bleeding related to the lock solution was encountered during the study period in either group."



Expert recommendations

The German guideline of infection prevention and hygiene and the US dialysis guideline (KDOQI) recommend urokinase for the prophylactic use in central venous catheters.


“
...Urokinase with or without taurolidine is a reasonable option for overproportional rates of bloodstream infections.
”¹²



TauroLock™ U25.000

TauroLock™-U25.000 contains the [antimicrobial](#) compound Taurolidine, 4% Citrate and Urokinase (25.000 IU).

TauroLock™-U25.000 fulfills the above mentioned guideline recommendations and contains Urokinase to ensure patency and provides infection control in the device.



Best practice & cost efficiency

Clinical outcomes		Sodium Citrate 4%	2+1 protocol TauroLock® HEP TAUROLOCK®	p value
Patency	catheter dysfunction per 1.000 catheter days	44.3	18.7	p = 0.001
	rt-PA Rate per 1.000 catheter days	9.8	3.8	p = 0.001
Bloodstream Infection	CRI per 1.000 catheter days	2.7	0.67	p = 0.003
Hospitalisation	CRI events in days	15.8	3.7	p = 0.001

Costs per year	Sodium Citrate 4%	2+1 protocol TauroLock® HEP TAUROLOCK®
costs of lock solution (purchase accumulated)	187.- USD	1.485.- USD
Costs for complication management (dysfunction, CRI)	4.309.- USD	1.063.- USD
Use of alteplase included!	→ 43 % cost savings	

Study authors: Winnicki, W. at al.¹³


Study location: Vienna, Austria

Patients enrolled: 106

Journal and publication date: Kidney International, 2017

Keynotes: "significant catheter-related cost savings of 43% in the taurolidine group vs. citrate group when overall expenses per patient and year were compared"

„use of taurolidine-based catheter lock solutions containing heparin and urokinase significantly reduced complications related to tunneled hemodialysis catheters when compared to four percent citrate solution and was overall more cost-efficient“



References

- ¹ Citrate versus heparin lock for hemodialysis catheters: a systematic review and meta-analysis of randomized controlled trials.
Zhao Y, Li Z, Zhang L, Yang J, Yang Y, Tang Y, Fu P. Am J Kidney Dis. 2014 Mar;63(3):479-90.
- ² Sodium citrate 4% locking solution for central venous dialysis catheters-an effective, more cost-efficient alternative to heparin.
Grudzinski L, Quinan P, Kwok S, Pierratos A. Nephrol Dial Transplant. 2007 Feb;22(2):471-6.
- ³ Citrate 4% versus Heparin and the Reduction of Thrombosis Study (CHARTS)
MacRae Jennifer M, Ivana Dojcinovic, Ognjenka Djurdjev, Beverly Jung, Steven Shalansky, Adeera Levin and Mercedeh Kiaii. CJASN March 2008, 3 (2) 369-374
- ⁴ Dialysis standard 2016, German Society of Nephrology in cooperation with the Association of German Kidney Centers as well as the Society of Paediatric Nephrology
- ⁵ Prophylaxis against Dialysis Catheter-Related Bacteremia with a Novel Antimicrobial Lock Solution.
Allon M, Clinical Infectious Diseases, 2003;36:1539-44
- ⁶ Observational study of need for thrombolytic therapy and incidence of bacteremia using taurolidine-citrate-heparin, taurolidine-citrate and heparin catheter locks in patients treated with hemodialysis.
Solomon LR, Cheesbrough JS, Bhargava R, Mitsides N, Heap M, Green G, Diggle P. Semin Dial 2012;25(2):233-238
- ⁷ Any use for alternative lock solutions in the prevention of catheter-related blood stream infections?
Labriola L, Pochet JM. J Vasc Access. 2017 Mar 6;18(Suppl. 1):34-38.
- ⁸ A new haemodialysis catheter-locking agent reduces infections in haemodialysis patients.
Taylor C, Cahill J, Gerrish M, Little J. J Ren Care 2008;34(3):116-120.
- ⁹ Tunneled catheters with taurolidine-citrate-heparin lock solution significantly improve the inflammatory profile of hemodialysis patients.
Fontseré N, Cardozo C, Donate J, Soriano A, Muros M, Pons M, Mensa J, Campistol JM, Navarro-Gonzalez JF, Maduell F. Antimicrob Agents Chemother 2014;58(7):4180-4184.
- ¹⁰ Taurolidine-citrate-heparin catheter lock solution reduces staphylococcal bacteraemia rates in haemodialysis patients.
Murray EC, Deighan C, Geddes C, Thomson PC. QJM 2014;107(12):995-1000.
- ¹¹ Safety and efficacy of taurolidine/urokinase versus taurolidine/heparin as a tunneled catheter lock solution in hemodialysis patients: a prospective, randomized, controlled study.
Al-Ali F, Hamdy AF, Hamad A, Elsayed M, Iqbal ZZ, Elsayed A, Ibrahim R, Tolba H, Buanan H, Fawzy A. NDT 2017:1-7. Guideline for infection prevention
- ¹² Guideline for infection prevention and hygiene 2019 in addition to the dialysis standard.
German society for nephrology e.V. 27.01.2020
- ¹³ Taurolidine-based catheter lock regimen significantly reduces overall costs, infection, and dysfunction rates of tunneled hemodialysis catheters.
Winnicki W, Herkner H, Lorenz M, Handisurya A, Kikic Z, Bielesz B, Schairer B, Reiter T, Eskandary F, Sunder-Plassmann G, Sengelge G. Kidney Int 2017; S0085-2538(17)30495-7.

